

Chicago Flu Update



City of Chicago Rahm Emanuel, Mayor

November 27, 2015

Chicago Department of Public Health Julie Morita, MD, Commissioner

What is the risk?

Currently, the risk of influenza infection is low. Vaccination is the best way to protect against influenza infection and all Chicagoans aged six months and older are encouraged to get vaccinated. Chicagoans should ask their healthcare provider or pharmacist about vaccine availability. For those without a healthcare provider or whose healthcare providers do not have the influenza vaccine, a schedule of <u>City of Chicago influenza vaccine clinics</u>¹ is available on the City website and by calling 311. To locate the closest City of Chicago clinic or retail pharmacy, go to <u>www.chicagoflushots.org</u>.

Are severe cases of influenza occurring?

The Illinois Department of Public Health (IDPH) has issued influenza testing and reporting recommendations². Suspected novel and variant influenza, pediatric influenza-associated deaths, influenza-associated ICU hospitalizations and outbreaks of influenza-like illness in a congregate setting should all be reported to CDPH via INEDSS³. For the week of November 15-21, 2015 (week 46), no influenza-associated ICU hospitalizations were reported (Figure 1).

Since October 4, 2015, five influenza-associated ICU hospitalizations have been reported; all were positive for influenza A (1 H3N2, 1 (H1N1)pdm09 and 3 unknown subtype [subtyping not attempted or not all subtypes tested]). Demographic and other aggregate information will be summarized once ten or more ICU hospitalizations have been reported.

How much influenza-like illness is occurring?

CDPH receives data from influenza surveillance sites across Chicago, which report the total number of patient visits seen weekly, and of those visits, the number with influenza-like illness (ILI) (i.e., fever of 100°F or greater, with cough or sore throat). All hospitals in Chicago that provide emergent care are required to report on a weekly basis the total number of emergency department visits, and of those visits, the number with ILI. For the week of November 15-21, 2015 with 11 hospitals reporting, 2.4% of emergency department visits were due to ILI (**Figure 2**).

ESSENCE is an electronic syndromic surveillance system that utilizes emergency department chief com-

Figure 1. Total number of reported influenza-associated ICU hospitalizations reported for Chicago residents, for current season (2015-2016) and previous season (2014-2015), Chicago, October-May.





Figure 2. Percent of <u>emergency department</u> visits attributed to influenza-like illness based on manual reporting as determined by individual hospitals, Chicago, by week, for the current season (2015-2016) and previous three seasons, October-May.



plaint data submitted daily by participating Chicago hospitals. ILI activity is determined solely based on the patient's chief complaint and does not take into account the entire medical record, as the ILI activity reported in Figure 2 does.

Currently, 10 Chicago hospitals submit data to ESSENCE. For the week of November 15-21, 2015, 1.2% of emergency department visits were due to ILI (**Figure 3**). Several outpatient clinics throughout Chicago participate in CDC's Influenza-like Illness Surveillance Network (**ILINet**) by reporting on the number of patients with ILI seen weekly. From November 15-21, 2015, with 24 facilities reporting, 4.4% of visits were due to influenza-like illness (**Figure 4**).

¹ http://www.cityofchicago.org/city/en/depts/cdph/supp_info/influenza/2015-influenza-walk-in-clinic-schedule.html; ² http://www.dph.illinois.gov/sites/default/files/publications/publicationsohpinfluenza-testing-and-reporting-memo-09.2015.pdf; ³https://dph.partner.illinois.gov/

Which influenza strains are circulating?

Data on influenza virus test results are reported by Chicago laboratories performing influenza subtyping. For the week of November 15-21, 2015, with 5 laboratories reporting, 6 of the 424 (1.4%) specimens tested for influenza were positive for influenza A [3 (H1N1) pdm09 and 3 unknown subtype]. Since October 4, 2015, 23 of 3,051 (<1%) specimens tested for influenza have been positive; 21 typed as influenza A (3 H3N2, 7 (H1N1)pdm09, 11 unknown subtype [subtyping not attempted or not all subtypes tested]) and 2 was typed as influenza B (Figure 5). The cumulative percent of specimens testing positive for influenza is similar to previous seasons during the same time period (Table 1).

Table 1. Cumulative number of specimens tested and positive for influenza for the current season (2015-2016) and the previous five seasons, Chicago, Weeks 40-46.

WEEKS TO TO:			
Influenza Season	Cumulative Tested	Cumulative Positive	Cumulative % Positive
2010-2011	992	3	<.5%
2011-2012	1,080	2	<.5%
2012-2013	1,521	29	2%
2013-2014	2,247	13	<1%
2014-2015	2,402	25	1%
2015-2016	3,051	23	<1%

Where can I get more information?

The Centers for Disease Control and Prevention's FluView⁴ report provides national updates and trends related to the intensity of influenza activity across the United States, as well as detailed information on antiviral resistance, severity of illness, and other topics. Updates specific to Illinois⁵ and Suburban Cook Coun- Figure 5. Percent of specimens testing positive (by RT-PCR) for ty⁶ are also available online. Current and archived issues of the Chicago Flu Update can be found on the CDPH website section Current Flu Situation in Chicago⁷. In 2013, the Metropolitan Chicago Healthcare Council (MCHC) and CDPH released "Stop the Spread: A Health Care Guide to Influenza Preparedness^{3,8}. This report provides an overview of influenza, it's impact on public health and how hospitals can prepare for, mitigate the impact of and respond to influenza infections and outbreaks.



Figure 3. Percent of emergency department visits attributed to influenza-like illness based on chief complaint data submitted to ESSENCE, Chicago, by week, for the current season (2015-2016) and the previous three seasons, October-May.






Figure 4. Percent of medically attended outpatient visits attributed to influenza-like illness as reported by ILINet facilities, Chicago, by week, for the current season (2015-2016) and the previous three seasons, October-May.



influenza by subtype as reported by local laboratories serving Chicago hospitals, for the current season (2015-2016) October-May.



All data are preliminary and may change as more reports are received.